

PSYCHOSIS

Psychosis refers to a loss of contact with reality. When people can't tell the difference between what is real and what is not, it is called a psychotic episode. A first episode of psychosis is often very frightening, confusing and distressing, particularly because it is an unfamiliar experience.

Psychosis usually first appears in a person's late teens or early 20s. Approximately three out of every 100 people will have a psychotic episode in their lifetime. Psychosis occurs in men and women and across all cultures and socioeconomic groups.

Psychotic illnesses seem to affect women at a later age than men, when women may be farther along in their social and work lives. On the whole, women respond better than men to most treatments. However, women have times when the risk of relapse is greater. These times are before their period is due, after childbirth and around menopause. This suggests that women's hormones may in some way affect psychosis.

A number of mental illnesses can include psychosis as a symptom, including:

- **Schizophrenia:** A person has some psychotic symptoms for at least six months, with a significant decline in the person's ability to function.
- **Schizophreniform disorder:** A person has some psychotic symptoms for less than six months.
- **Bipolar disorder:** With this type of illness, the symptoms of psychosis relate more to mood disturbance than to thought disturbance.
- **Schizoaffective disorder:** A person will have symptoms of schizophrenia and symptoms of a mood disturbance, either at the same time or alternating over time.
- **Depression with psychotic features:** A person has severe depression and symptoms of psychosis without the mania associated with bipolar disorder.
- **Drug-induced psychosis:** The use of drugs such as marijuana, cocaine, LSD, amphetamines and alcohol can sometimes cause psychotic symptoms to appear.
- **Organic psychosis:** Symptoms of psychosis may appear as a result of a physical illness or a head injury.
- **Brief psychotic disorder:** This illness usually lasts less than a month. It is usually triggered by a major stress in the person's life, such as a death in the family.
- **Delusional disorder:** This type of psychosis consists of very strong and fixed beliefs in things that are not true.
- **Post-traumatic stress disorder:** This usually lasts more than a month, and happens after a person has seen or experienced a very traumatic event. The person may have flashbacks or hallucinate.

SYMPTOMS

Psychosis can come on suddenly or can develop very gradually. Symptoms of psychosis can vary from person to person and may change over time. Some common symptoms are:

- changes in thinking patterns (difficulty concentrating; loss of memory; disconnected thoughts)
- delusions (fixed, false beliefs that are not consistent with the person's culture, and have no basis in fact)
- hallucinations (people hear, see, taste, smell or feel something that does not actually exist)
- changes in mood (finding it hard to express feelings; feeling inappropriate or intense bursts of emotion; feeling empty of any emotions; depression)
- very disorganized behaviour (cannot complete everyday tasks such as bathing, dressing appropriately and preparing simple meals)
- thoughts of death or suicide.

CAUSES

In most cases, we do not know what causes a first episode of psychosis. Current research shows that biological and genetic factors raise the risk of having psychosis. Brain chemistry may affect psychosis.

A person who is having symptoms of psychosis should have a thorough medical assessment, to rule out any physical illness that may be the cause.

TREATMENTS

Psychosis can be treated, and many people make a good recovery, especially if they get help early. Treatment may be either on an out-patient basis or in hospital. Treatment usually consists of medication and psychosocial interventions (counselling, for example).

Throughout treatment, families can receive support and education during sessions with the treatment team.

Medication called antipsychotic medication is usually essential. It relieves symptoms of psychosis and may prevent further episodes of illness.

A case manager or therapist can provide emotional support, education about the illness and its management, and practical assistance with day-to-day living. They may also recommend programs in the community and provide supportive psychotherapy and vocational counselling.

RECOVERY

Recovery from a first episode of psychosis varies from person to person. Sometimes symptoms go away quickly and people are able to resume their regular life right away. Other people may need several weeks or months to recover. Some people will need medication and support for the rest of their lives.



For further information:

R. Samuel McLaughlin Addiction and Mental
Health Information Centre
1 800 463-6273 or 416 595-6111 in Toronto

Visit our Web site at www.camh.net

For information on other Centre for Addiction
and Mental Health resource materials or to place
an order, please contact:

Marketing and Sales Services
33 Russell St., Toronto, Ontario M5S 2S1
Tel: 1 800 661-1111 or 416 595-6059 in Toronto
Email: marketing@camh.net

Adapted from *First Episode Psychosis: A Guide for
People with Psychosis and their Families* © 1999,
Centre for Addiction and Mental Health